

Product label

TECH FLO[®] CAL-BOR+MOLY
10.0% Calcium, 8.0% Sulfur, 1.0% Boron, 0.5% Molybdenum

TECH-FLO CAL-BOR+MOLY is a neutral, micronized powder formulated as a flowable dispersion for use as a foliar spray.

GUARANTEED ANALYSIS

Calcium (Ca).....10.0%
Sulfur(S)..... 8.0%
Boron (B).....1.0%
Molybdenum (Mo).....0.5%

calcium and Sulfur derived from hydrated calcium sulfate. Boron derived from boric acid and sodium borate. Molybdenum derived from ammonium molybdate.

GENERAL RECOMMENDATIONS

AS A NUTRITIONAL SPRAY: SEE SUGGESTED USES

DILUTION: Apply the suggested rates of TECH-FLO CAL-BOR+MOLY in the following amounts of water per acre:

Field & Row Crops: By air: 3-20 gals. By ground: 10 gals. minimum

Tree Crops: By air: 10 gals. minimum. By ground concentrate: 40-100 gals. By ground dilute:100-500 gals.

In oil sprays, use 1-4 pints per 100 gallons

COMPATIBILITY: This product has been designed to be compatible with most commonly used insecticides, fungicides and acaricides. However, it is not feasible to test all possible combinations of pesticides with which it may be mixed. A compatibility test is recommended for combinations that have not been used previously. Consult your NUTRI-TECH representative for detailed compatibility information

WARNING: This product contains Boron and is to be used only where Boron is known to be beneficial and on the crops recommended on this label. Any other use may result in injury to the crops.

WARNING: Crops high in molybdenum are toxic to grazing animals (ruminants).

CAUTION: Causes skin irritation. Avoid contact with skin, eyes, open cuts or sores. In case of contact, flush with water. Get medical attention for eyes.

CAUTION
KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED

This product has been registered with the Washington State Department of Agriculture. When applied as directed, this fertilizer meets the Washington standards for arsenic, cadmium, cobalt, mercury, molybdenum, lead, nickel, selenium and zinc. You have the right to receive specific information about Washington standards from the distributor of this product.

**SHAKE WELL BEFORE USING
ADEQUATE AGITATION SHOULD BE MAINTAINED AT ALL TIMES DURING MIXING AND
APPLICATION**

SUGGESTED USES:

Alfalfa, Hay & Other Forage Crops - Apply 1-4 quarts over spring growth, repeat after each cutting.

Almonds - Apply 1-4 quarts at 2-3 week intervals starting pre-bloom.

Apples, Pears - Apply 2-4 quarts at 2-4 week intervals starting pre-bloom.

Barley, Oats, Rice, Wheat & Other Small Grains - Apply 1-4 quarts at tillering, repeat at early boot if necessary.

Beans, Peas, & related Legumes - Apply 1-4 quarts at early bud, repeat after bloom if necessary.

Cabbage, Cauliflower, Broccoli, Brussels Sprouts & Other Cole Crops - Apply 1-4 quarts as needed after thinning or transplant.

Carrots, Onions, Radishes, Beets & Other Vegetable Root Crops - Apply 1-4 quarts when top growth is 4-8" high, repeat as necessary.

Citrus, Avocados - Apply 2-4 quarts in spring, summer, and fall sprays.

Corn, Field & Sweet - Apply 1-4 quarts when 10-20" high, repeat in 2-4 weeks as necessary.

Cotton - Apply 1-4 quarts at first square, repeat in 2-4 weeks as necessary.

Cucumbers, Melons, Squash & Other Cucurbits - Apply 1-4 quarts prebloom, repeat in 2-4 weeks as necessary.

Grapes - Apply 1-4 quarts dormant or as needed pre-bloom to post-bloom.

Hops - Apply 1-4 quarts as needed in cover sprays.

Lettuce, Celery, Spinach & Other Green Leafy Vegetables - Apply 1-4 quarts as needed after thinning and transplant.

Mint - Apply 1-4 quarts over new growth, repeat as needed.

Peaches, Plums, Cherries & Other Stone Fruits - Apply 1-4 quarts pre-bloom and at 3-4 week intervals as needed post-bloom

Peanuts - Apply 1-4 quarts at early bloom and just prior to pegging.

Pecans & Other Deciduous Nut Trees - Apply 1-4 quarts at 3-4 week intervals starting at bud break.

Ornamentals, Trees & Shrubs - Apply 1-4 quarts at 4-6 week intervals as needed.

Potatoes - Apply 1-4 quarts at tuber initiation, repeat in 2-4 weeks as necessary.

Raspberries, Blackberries, Blueberries & Other Small Fruit - Apply 1-4 quarts pre-bloom, repeat in 2-4 weeks as necessary.

Soybeans, Seed Crops - Apply 1-4 quarts pre-bloom, repeat in 2-4 weeks as necessary.

Sugarbeets - Apply 1-4 quarts at 4th-6th leaf, repeat in 2-4 weeks as necessary.

Strawberries - Apply 1-4 quarts pre-bloom, repeat in 2-4 weeks as necessary.

Tomatoes, Peppers, Eggplant & Okra - Apply 1-4 quarts pre-bloom, repeat in 2-4 weeks as necessary.

Turf (Farms) - Apply 1-4 quarts at 4-6 week intervals as needed.

Walnuts - Apply 1-4 quarts in blight sprays.

NOTICE: Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used according to directions under normal conditions on the crops specified. Because of widely varying use conditions, it is impossible to eliminate all risks even when the label directions are followed. Therefore, the manufacturer makes no other warranties, expressed or implied. The buyer assumes all risks associated with the use of this product. In the event of damage resulting from a breach of warranty, the buyer agrees to accept rotund of the purchase price as full discharge of the manufacturer's liability.

**NUTRIENT TECHNOLOGIES, INC
DINUBA, CA 93618**

**NET CONTENTS: 5 GALLONS (18.925 LITERS)
NET WT 57.0 LB.**

©1993. Nutrient Technologies, Inc.

[*Return to Home Page*](#)

[*Return to Selection Chart*](#)
